

WHEELCHAIR CPR



Ensure Safety for Self & Others
Dial Triple Zero (000)
For an Ambulance

Preparing a casualty in a wheelchair for CPR

DANGER

Remove hazards

RESPONSE

Talk, touch, squeeze shoulders

SEND FOR HELP

Call triple zero (000)

AIRWAY

Open with head tilt/chin lift

BREATHING

Look, listen and feel

MAKE ROOM BEHIND THE CHAIR - APPLY THE BRAKES - REMOVE ACCESSORIES

(armrests, anti-tip devices, food tray foot plates and other removeable parts)

ONE FIRST AIDER

If possible, secure the casualty to the chair
(ie around the chest and chair back with a belt or similar)

Stand behind the chair

Hold on to the handles

TWO FIRST AIDERS

One on each side of the chair

Kneel down on one knee
(with the foot at the rear of the chair on the ground)

Place your arm under the casualty's armpit
and hold onto the handle

The other hand supports the casualty's head

MORE THAN TWO FIRST AIDERS

If several people can help:

Make room on the floor

Carefully lift the casualty
from the wheelchair

Place casualty on their
back on the ground

LOWER THE CHAIR BACKWARDS TO THE GROUND

REMOVE FROM THE CHAIR

Support under armpits, pull/slide out of the chair onto the ground

START CPR

30 chest compressions
2 breaths

SIGNS & SYMPTOMS

- Cannot be woken.
- Unresponsive.
- Not breathing normally or at all.
- Pale or blue in colour

IMPORTANT

Alternative methods may apply.

Follow care plan if available.

Use good manual handling techniques.

Do not attempt if beyond your strength capabilities.

Avoiding injury to the casualty and the first aider is the priority.

Support and protect the head.

Only do CPR in the chair if it has a hard back.

May not be suitable for motorised wheelchairs.

Do not do this in a plastic chair.